

$B_s \rightarrow (\rho, \omega, \phi)\eta^{(\prime)}$ Decays in the Perturbative QCD Approach

Xin-fen Chen,^{*} Dong-qin Guo,[†] and Zhen-jun Xiao[‡]

*Department of Physics and Institute of Theoretical Physics,
Nanjing Normal University, Nanjing, Jiangsu 210097, P.R.China*

(Dated: February 2, 2008)

Abstract

In this paper, we calculate the branching ratios and CP-violating asymmetries for $B_s \rightarrow (\rho^0, \omega, \phi)\eta^{(\prime)}$ decays in the perturbative QCD (pQCD) factorization approach. Numerically we found that (a) the pQCD predictions for the CP-averaged branching ratios are $Br(B_s \rightarrow \rho^0\eta) \approx 0.07 \times 10^{-6}$, $Br(B_s \rightarrow \rho^0\eta') \approx 0.10 \times 10^{-6}$, $Br(B_s \rightarrow \omega\eta) \approx 0.02 \times 10^{-6}$, $Br(B_s \rightarrow \omega\eta') \approx 0.13 \times 10^{-6}$, $Br(B_s \rightarrow \phi\eta) \approx 2.7 \times 10^{-5}$ and $Br(B_s \rightarrow \phi\eta') \approx 2.0 \times 10^{-5}$; (b) the gluonic contributions are small in size: less than 3% for $B \rightarrow (\rho, \omega, \phi)\eta$ decays, and about 10% for $B \rightarrow (\rho, \omega, \phi)\eta'$ decays; and (c) the pQCD predictions for the CP-violating asymmetries of the considered decays are generally not large in magnitude. The above predictions can be tested in the forthcoming LHC-b experiments at CERN.

PACS numbers: 13.25.Hw, 12.38.Bx, 14.40.Nd

^{*}Electronic address: chenxinfen@163.com

[†]Electronic address: medongqin@163.com

[‡]Electronic address: xiaozhenjun@njnu.edu.cn

I. INTRODUCTION

The theoretical calculations and experimental measurements of the rare B meson decays play an important role in testing the standard model (SM), probing CP violation of B meson system and searching for possible new physics beyond the SM. At present, about 1000 million events of B meson pair productions and decays have been collected by BaBar and Belle collaborations. In the forthcoming LHC experiments, a huge amount of B meson events, say around $10^{11} \sim 10^{12}$, are expected, and therefore the rare B meson decays with a branching ratio around 10^{-7} can be observed with good precision. Another advantage is that the heavier B_s and B_c mesons and b-baryons, besides the B_u and B_d , can also be produced and studied at LHC [1].

By employing the generalized factorization approach[2, 3] or the QCD factorization (QCDF) approach [4], about forty $B_s \rightarrow h_1 h_2$ (h_i stand for light pseudo-scalar or vector mesons) decay modes have been studied in the framework of SM [5, 6, 7, 8] or in some new physics models beyond the SM [9]. In this paper, we will study the $B_s \rightarrow \rho^0 \eta^{(\prime)}$, $\omega \eta^{(\prime)}$ and $\phi \eta^{(\prime)}$ decays in the pQCD factorization approach. In principle, the physics for the B_s two-body hadronic decays is very similar to that for the B_d meson except that the spectator d quark is replaced by the s quark.

For $B_s \rightarrow (\rho^0, \omega, \phi) \eta^{(\prime)}$ decays, the B_s meson is heavy, setting at rest and decaying into two light mesons (i.e. ρ^0 and $\eta^{(\prime)}$) with large momenta. Therefore the light final state mesons are moving very fast in the rest frame of B_s meson. In this case, the short distance hard process dominates the decay amplitude. We assume that the soft final state interaction is not important for such decays. The smallness of FSI effects for B meson decays into two light final state mesons has been put forward by Bjorken [10] based on the color transparency argument [11], and also supported by further renormalization group analysis of soft gluon exchanges among initial and final state mesons [12]. With the Sudakov resummation, we can include the leading double logarithms for all loop diagrams, in association with the soft contribution. Unlike the usual factorization approach, the hard part of the pQCD approach consists of six quarks rather than four. We thus call it six-quark operators or six-quark effective theory. Applying the six-quark effective theory to B_s meson decays, we need meson wave functions for the hadronization of quarks into mesons. All the collinear dynamics are included in the meson wave functions.

This paper is organized as follows. In Sec. II, we give a brief review for the PQCD factorization approach. In Sec. III, we calculate analytically the related Feynman diagrams and present the decay amplitudes for the studied decay modes. In Sec. IV, we show the numerical results for the branching ratios and CP asymmetries of $B_s \rightarrow \rho^0 \eta^{(\prime)}$, $\omega \eta^{(\prime)}$ and $\phi \eta^{(\prime)}$ decays and comparing them with the results obtained in the other two methods mentioned above. The summary and some discussions are included in the final section.

II. THEORETICAL FRAMEWORK

The pQCD factorization approach has been developed and applied in the non-leptonic B meson decays for some time [11, 13, 14, 15]. This approach is based on k_T factorization scheme, where three energy scales are involved [13]. In this approach, the decay amplitude is factorized into the convolution of the meson's light-cone wave functions, the

hard scattering kernels and the Wilson coefficients, which stand for the soft (Φ), hard(H), and harder(C) dynamics respectively. The hard dynamics (H) describes the four quark operator and the spectator quark connected by a hard gluon. This hard part is characterized by $\sqrt{\Lambda M_{B_s}}$, and can be calculated perturbatively in pQCD approach. The harder dynamics (C) is from m_W scale to m_{B_s} scale described by renormalization group equation for the four quark operators. The dynamics below $\sqrt{\Lambda M_{B_s}}$ is soft, which is described by the meson wave functions (Φ). While the function (H) depends on the processes considered, the wave function is independent of the specific processes. Using the wave functions determined from other well measured processes, one can make quantitative predictions here. Based on this factorization, the decay amplitude can be written as the following

$$\mathcal{A}(B_s \rightarrow M_1 M_2) \sim \int d^4 k_1 d^4 k_2 d^4 k_3 \text{Tr} [C(t) \Phi_{B_s}(k_1) \Phi_{M_1}(k_2) \Phi_{M_2}(k_3) H(k_1, k_2, k_3, t)], \quad (1)$$

where k_i 's are momenta of light quarks included in each mesons, and Tr denotes the trace over Dirac and color indices. $C(t)$ is Wilson coefficient of the four quark operator which results from the radiative corrections at short distance. The functions Φ_M and H are meson wave functions and the hard part respectively.

For the B_s meson decays, since the b quark is rather heavy we consider the B_s meson at rest for simplicity. It is convenient to use light-cone coordinate $(p^+, p^-, \mathbf{p}_T)$ to describe the meson's momenta,

$$p^\pm = \frac{1}{\sqrt{2}}(p^0 \pm p^3), \quad \text{and} \quad \mathbf{p}_T = (p^1, p^2). \quad (2)$$

Using these coordinates the B_s meson and the two final state meson momenta can be written as:

$$P_{B_s} = \frac{M_{B_s}}{\sqrt{2}}(1, 1, \mathbf{0}_T), \quad P_\rho = \frac{M_{B_s}}{\sqrt{2}}(1, r_\rho^2, \mathbf{0}_T), \quad P_\eta = \frac{M_{B_s}}{\sqrt{2}}(0, 1 - r_\rho^2, \mathbf{0}_T), \quad (3)$$

respectively, where $r_\rho = m_\rho/M_{B_s}$ and the light pseudoscalar meson masses have been neglected (here we just take the decay $B_s \rightarrow \rho^0 \eta$ as an example). Putting the light (anti-) quark momenta in B_s , ρ^0 and η mesons as k_1 , k_2 , and k_3 , respectively, we can choose

$$k_1 = (x_1 P_1^+, 0, \mathbf{k}_{1T}), \quad k_2 = (x_2 P_2^+, 0, \mathbf{k}_{2T}), \quad k_3 = (0, x_3 P_3^-, \mathbf{k}_{3T}). \quad (4)$$

Then the integration over k_1^- , k_2^- , and k_3^+ in eq.(1) will lead to

$$\mathcal{A}(B_s \rightarrow \rho^0 \eta) \sim \int dx_1 dx_2 dx_3 b_1 db_1 b_2 db_2 b_3 db_3 \cdot \text{Tr} [C(t) \Phi_{B_s}(x_1, b_1) \Phi_\rho(x_2, b_2) \Phi_\eta(x_3, b_3) H(x_i, b_i, t) S_t(x_i) e^{-S(t)}], \quad (5)$$

where b_i is the conjugate space coordinate of k_{iT} , and t is the largest energy scale in function $H(x_i, b_i, t)$, as a function in terms of x_i and b_i . The large logarithms ($\ln m_W/t$) coming from QCD radiative corrections to four quark operators are included in the Wilson coefficients $C(t)$. The large double logarithms ($\ln^2 x_i$) on the longitudinal direction are summed by the threshold resummation [15], and they lead to $S_t(x_i)$ which smears the end-point singularities on x_i . The last term, $e^{-S(t)}$, is the Sudakov form factor resulting from

overlap of soft and collinear divergences, which suppresses the soft dynamics effectively. Thus it makes the perturbative calculation of the hard part H applicable at intermediate scale, i.e., M_{B_s} scale. We will calculate analytically the function $H(x_i, b_i, t)$ for our six decays in the first order in α_s expansion and give the convoluted amplitudes in next section.

For the considered $B_s \rightarrow \rho^0 \eta^{(\prime)}, \omega \eta^{(\prime)}$ and $\phi \eta^{(\prime)}$ decays, the low energy weak effective Hamiltonian $\mathcal{H}_{eff}$ for the $b \rightarrow q$ transition with $q = (d, s)$ can be written as [16]:

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \left[V_{ub} V_{uq}^* (C_1(\mu) O_1^u(\mu) + C_2(\mu) O_2^u(\mu)) - V_{tb} V_{tq}^* \sum_{i=3}^{10} C_i(\mu) O_i(\mu) \right]. \quad (6)$$

We specify below the operators in $\mathcal{H}_{eff}$ for $b \rightarrow d$ transition:

$$\begin{aligned} O_1^u &= \bar{d}_\alpha \gamma^\mu L u_\beta \cdot \bar{u}_\beta \gamma_\mu L b_\alpha, & O_2^u &= \bar{d}_\alpha \gamma^\mu L u_\alpha \cdot \bar{u}_\beta \gamma_\mu L b_\beta, \\ O_3 &= \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu L q'_\beta, & O_4 &= \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu L q'_\alpha, \\ O_5 &= \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu R q'_\beta, & O_6 &= \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu R q'_\alpha, \\ O_7 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu R q'_\beta, & O_8 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu R q'_\alpha, \\ O_9 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu L q'_\beta, & O_{10} &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu L q'_\alpha, \end{aligned} \quad (7)$$

where α and β are the $SU(3)$ color indices; $L, R = (1 \mp \gamma_5)$ are the left- and right-handed projection operators. The sum over q' runs over the quark fields that are active at the scale $\mu = O(m_b)$. Since we here work at the leading twist approximation and leading double logarithm summation, we will also use the leading order (LO) expressions of the Wilson coefficients $C_i(\mu)$ ($i = 1, \dots, 10$), although the next-to-leading order $C_i(\mu)$ already exist in the literature [16]. This is the consistent way to cancel the explicit μ dependence in the theoretical formulae.

For the renormalization group evolution of the Wilson coefficients from higher scale to lower scale, we use the formulae as given in Ref. [14] directly. At the high m_W scale, the leading order Wilson coefficients $C_i(M_W)$ are simple and can be found easily in Ref. [16]. In pQCD approach, the scale ' t ' may be larger or smaller than the m_b scale. For the case of $m_b < t < m_W$, we evaluate the Wilson coefficients at t scale using leading logarithm running equations, as given in Eq.(C1) of Ref. [14]. For the case of $t < m_b$, we then evaluate the Wilson coefficients at t scale by using $C_i(m_b)$ as input and the formulae given in Appendix D of Ref. [14].

For the wave function of the heavy B_s meson, we take

$$\Phi_{B_s} = \frac{1}{\sqrt{2N_c}} (\not{p}_{B_s} + M_{B_s}) \gamma_5 \phi_{B_s}(\mathbf{k}_1). \quad (8)$$

Here only the contribution of Lorentz structure $\phi_{B_s}(\mathbf{k}_1)$ is taken into account, since the contribution of the second Lorentz structure $\tilde{\phi}_{B_s}$ is numerically small and can be neglected. The distribution amplitude ϕ_{B_s} in Eq. (8) will be given lately in Eq. (47).

For $B \rightarrow V \eta^{(\prime)}$ decays, the vector meson $V = (\rho, \phi, \omega)$ is longitudinally polarized. The relevant longitudinal polarized component of the wave function for ρ meson, for example, is given as [17],

$$\Phi_\rho = \frac{1}{\sqrt{2N_c}} \{ \not{\epsilon} [m_\rho \phi_\rho(x) + \not{p}_\rho \phi_\rho^t(x)] + m_\rho \phi_\rho^s(x) \}, \quad (9)$$

where the first term is the leading twist wave function (twist-2), while the second and third terms are sub-leading twist (twist-3) wave functions. For the case of $V = \omega$ and ϕ , their wave functions are the same in structure as that defined in Eq. (9), but with different distribution amplitudes. One can find the distribution amplitudes $\phi_{\omega,\phi}$ and $\phi_{\omega,\phi}^{t,s}(x)$ in next section.

For η and η' meson, the wave function for $d\bar{d}$ components of $\eta^{(\prime)}$ meson are given as [18]

$$\Phi_{\eta_{d\bar{d}}}(P, x, \zeta) \equiv \frac{i\gamma_5}{\sqrt{2N_c}} \left[\not{p} \phi_{\eta_{d\bar{d}}}^A(x) + m_0^{\eta_{d\bar{d}}} \phi_{\eta_{d\bar{d}}}^P(x) + \zeta m_0^{\eta_{d\bar{d}}} (\not{p} \not{\gamma} - v \cdot n) \phi_{\eta_{d\bar{d}}}^T(x) \right], \quad (10)$$

where P and x are the momentum and the momentum fraction of $\eta_{d\bar{d}}$ respectively, while $\phi_{\eta_{d\bar{d}}}^A$, $\phi_{\eta_{d\bar{d}}}^P$ and $\phi_{\eta_{d\bar{d}}}^T$ represent the axial vector, pseudoscalar and tensor components of the wave function respectively, and will be given explicitly in next section. Following Ref. [18], we here also assume that the wave function of $\eta_{d\bar{d}}$ is the same as π wave function based on SU(3) flavor symmetry. The parameter ζ is either +1 or -1 depending on the assignment of the momentum fraction x .

Before we proceed to do the perturbative calculations, we firstly give a brief discussion about the $\phi - \omega$ mixing, as well as the $\eta - \eta'$ mixing and the gluonic component of the $\eta^{(\prime)}$ mesons.

For the vector $\phi - \omega$ meson system, we choose the ideal mixing scheme between $\phi(1020)$ and $\omega(782)$

$$\omega = \frac{1}{\sqrt{2}} (u\bar{u} + d\bar{d}), \quad \phi = -s\bar{s}, \quad (11)$$

since the current data support this ideal mixing scheme [19]. The quark contents of ρ^0 meson is chosen as $\rho^0 = (-u\bar{u} + d\bar{d})/\sqrt{2}$ [19].

For the $\eta - \eta'$ system, there exist two popular mixing basis: the octet-singlet basis and the quark-flavor basis [20, 21]. Here we use the quark-flavor basis [20] and define

$$\eta_q = (u\bar{u} + d\bar{d})/\sqrt{2}, \quad \eta_s = s\bar{s}. \quad (12)$$

The physical states η and η' are related to η_q and η_s through a single mixing angle ϕ ,

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = U(\phi) \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix}. \quad (13)$$

The corresponding decay constants $f_q, f_s, f_\eta^{q,s}$ and $f_{\eta^{(\prime)}}^{q,s}$ have been defined in Ref. [20] as

$$\begin{aligned} \langle 0 | \bar{q} \gamma^\mu \gamma_5 q | \eta_q(P) \rangle &= -\frac{i}{\sqrt{2}} f_q P^\mu, \\ \langle 0 | \bar{s} \gamma^\mu \gamma_5 s | \eta_s(P) \rangle &= -i f_s P^\mu, \end{aligned} \quad (14)$$

$$\begin{aligned} \langle 0 | \bar{q} \gamma^\mu \gamma_5 q | \eta^{(\prime)}(P) \rangle &= -\frac{i}{\sqrt{2}} f_{\eta^{(\prime)}}^q P^\mu, \\ \langle 0 | \bar{s} \gamma^\mu \gamma_5 s | \eta^{(\prime)}(P) \rangle &= -i f_{\eta^{(\prime)}}^s P^\mu, \end{aligned} \quad (15)$$

while the decay constants $f_\eta^{q,s}$ and $f_{\eta'}^{q,s}$ are related to f_q and f_s via the same mixing matrix,

$$\begin{pmatrix} f_\eta^q & f_\eta^s \\ f_{\eta'}^q & f_{\eta'}^s \end{pmatrix} = U(\phi) \begin{pmatrix} f_q & 0 \\ 0 & f_s \end{pmatrix}. \quad (16)$$

The three input parameters f_q , f_s and ϕ in the quark-flavor basis have been extracted from various related experiments [20, 21]

$$f_q = (1.07 \pm 0.02)f_\pi, \quad f_s = (1.34 \pm 0.06)f_\pi, \quad \phi = 39.3^\circ \pm 1.0^\circ, \quad (17)$$

where $f_\pi = 130$ MeV. In the numerical calculations, we will use these mixing parameters as inputs.

As shown in Eq. (13), the physical states η and η' are generally considered as a linear combination of light quark pairs $u\bar{u}$, $d\bar{d}$ and $s\bar{s}$. But it should be noted that the η' meson may have a gluonic component. Following Ref. [22], we also estimated the possible gluonic contributions to $B \rightarrow (\rho, \omega, \phi)\eta^{(\prime)}$ decays induced by the gluonic corrections to the $B \rightarrow \eta^{(\prime)}$ transition form factors [22] and found that these corrections to both the branching ratios and CP violating asymmetries are indeed small: less than 10%.

Frankly speaking, on the other hand, we currently still do not know how to calculate reliably the gluonic contributions to the B meson decays involving η' meson as final state particle. For the studied decay modes in this paper, we firstly consider only the dominant contributions from the quark contents of $\eta^{(\prime)}$ meson, and then take the subdominant contribution from the possible gluonic content of $\eta^{(\prime)}$ meson as a source of theoretical uncertainties.

III. PERTURBATIVE CALCULATIONS

In this section, we will calculate the hard part $H(t)$ for the considered decays. Following the same procedure as being used in Refs. [23, 24, 25, 26] and taking $B \rightarrow \rho^0\eta^{(\prime)}$ decays as an example, we will calculate and show the analytical results for all decay amplitudes, and then extend the results of $B \rightarrow \rho^0\eta^{(\prime)}$ decays to other decay modes under study.

As illustrated in Fig. 1, there are eight type Feynman diagrams contributing to the $B_s \rightarrow \rho^0\eta^{(\prime)}$ decays. We first calculate the usual factorizable diagrams (a) and (b). Operators O_1 , O_2 , O_3 , O_4 , O_9 , and O_{10} are $(V-A)(V-A)$ currents, the sum of their amplitudes can be written as:

$$\begin{aligned} F_{e\eta} = & -8\pi C_F M_{B_s}^A f_\rho \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \phi_{B_s}(x_1, b_1) \\ & \cdot \left\{ [(1+x_3)\phi_\eta^A(x_3, b_3) + r_\eta(1-2x_3)(\phi_\eta^P(x_3, b_3) + \phi_\eta^T(x_3, b_3))] \right. \\ & \alpha_s(t_e^1) h_e(x_1, x_3, b_1, b_3) \exp[-S_{ab}(t_e^1)] \\ & \left. + 2r_\eta \phi_\eta^P(x_3, b_3) \alpha_s(t_e^2) h_e(x_3, x_1, b_3, b_1) \exp[-S_{ab}(t_e^2)] \right\}. \end{aligned} \quad (18)$$

where $C_F = 4/3$ is a color factor and $r_\eta = m_0^{\eta s\bar{s}}/M_{B_s}$ or $r_\eta = m_0^{\eta d\bar{d}}/M_{B_s}$ since r_η depends on the quark components in η . The function h_e^i , the scales t_e^i and the Sudakov factors S_{ab} are displayed in the appendix. From diagrams Fig. 1(a) and 1(b), one can also extract out the form factor $F_0^{B_s \rightarrow \eta s\bar{s}}$.




FIG. 1: Diagrams contributing to the $B_s \rightarrow \rho^0 \eta$ decay (diagram (a) and (b) contribute to the $B_s \rightarrow \eta_s \bar{s}$ form factor $F_0^{B_s \rightarrow \eta_s \bar{s}}$).

The operators O_5 , O_6 , O_7 , and O_8 have a structure of $(V - A)(V + A)$. In some decay channels, some of these operators contribute to the decay amplitude in a factorizable way. Since only the vector part of $(V \pm A)$ current contribute to the scalar meson production,

$$\langle \rho | V + A | B \rangle \langle \eta | V - A | 0 \rangle = \langle \rho | V - A | B \rangle \langle \eta | V - A | 0 \rangle, \quad (19)$$

the result of these operators is the same as Eq. (18), and therefore we find easily that

$$F_{e\eta}^{P_1} = F_{e\eta}. \quad (20)$$

In some other cases, one needs to do Fierz transformation for these operators to get right flavor and color structure for factorization to work. In this case, we get $(S + P)(S - P)$ operators from $(V - A)(V + A)$ ones. For these $(S + P)(S - P)$ operators, Fig. 1(a) and 1(b) will give

$$F_{e\eta}^{P_2} = 0. \quad (21)$$

For the non-factorizable diagrams 1(c) and 1(d), all three meson wave functions are involved. The integration of b_3 can be performed by using $\delta(b_3 - b_1)$, leaving

only integration of b_1 and b_2 . For the $(V - A)(V - A)$ operators, the corresponding decay amplitude is:

$$M_{e\eta} = \frac{32}{\sqrt{6}}\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \cdot \phi_{B_s}(x_1, b_1) \phi_\rho(x_2, b_2) \{x_3 [\phi_\eta^A(x_3, b_1) - 2r_\eta \phi_\eta^T(x_3, b_1)] \cdot \alpha_s(t_f) h_f(x_1, x_2, x_3, b_1, b_2) \exp[-S_{cd}(t_f)]\} . \quad (22)$$

For the $(S - A)(V + A)$ operators, the decay amplitudes read

$$M_{e\eta}^{P_1} = \frac{64}{\sqrt{6}}\pi C_F M_{B_s}^4 r_\rho \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_{B_s}(x_1, b_1) \cdot \{ [x_2 \phi_\eta^A(x_3, b_1) (\phi_\rho^s(x_2, b_2) - \phi_\rho^t(x_2, b_2)) + r_\eta ((x_2 + x_3) (\phi_\eta^P(x_3, b_1) \cdot \phi_\rho^s(x_2, b_2) + \phi_\eta^T(x_3, b_1) \phi_\rho^t(x_2, b_2)) + (x_3 - x_2) (\phi_\eta^P(x_3, b_1) \phi_\rho^t(x_2, b_2) + \phi_\eta^T(x_3, b_1) \phi_\rho^s(x_2, b_2)))] \alpha_s(t_f) h_f(x_1, x_2, x_3, b_1, b_2) \exp[-S_{cd}(t_f)] \} , \quad (23)$$

$$M_{e\eta}^{P_2} = -M_{e\eta} . \quad (24)$$

For the non-factorizable annihilation diagrams 1(e) and 1(f), again all three wave functions are involved. Here we have all three kinds of contributions. $M_{a\eta}$ is the contribution containing operator type $(V - A)(V - A)$, while $M_{a\eta}^{P_1}$ and $M_{a\eta}^{P_2}$ is the contribution containing operator type $(V - A)(V + A)$ and $(S - P)(S + P)$, respectively

$$M_{a\eta} = -\frac{32}{\sqrt{6}}\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_{B_s}(x_1, b_1) \cdot \{ - [x_2 \phi_\rho(x_2, b_2) \phi_\eta^A(x_3, b_2) + r_\rho r_\eta ((x_2 - x_3) (\phi_\eta^P(x_3, b_2) \phi_\rho^t(x_2, b_2) + \phi_\eta^T(x_3, b_2) \phi_\rho^s(x_2, b_2)) + (2 + x_2 + x_3) \phi_\eta^P(x_3, b_2) \phi_\rho^s(x_2, b_2) + (-2 + x_2 + x_3) \phi_\eta^T(x_3, b_2) \phi_\rho^t(x_2, b_2))] \alpha_s(t_f^2) h_f^2(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^2)] + [x_3 \phi_\rho(x_2, b_2) \phi_\eta^A(x_3, b_2) + r_\rho r_\eta ((x_3 - x_2) (\phi_\eta^P(x_3, b_2) \phi_\rho^t(x_2, b_2) + \phi_\eta^T(x_3, b_2) \phi_\rho^s(x_2, b_2)) + (x_2 + x_3) (\phi_\eta^P(x_3, b_2) \phi_\rho^s(x_2, b_2) + \phi_\eta^T(x_3, b_2) \phi_\rho^t(x_2, b_2)))] \alpha_s(t_f^1) h_f^1(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^1)] \} . \quad (25)$$

$$M_{a\eta}^{P_1} = -\frac{32}{\sqrt{6}}\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_{B_s}(x_1, b_1) \cdot \{ [r_\rho (2 - x_2) \phi_\eta^A(x_3, b_2) (\phi_\rho^t(x_2, b_2) + \phi_\rho^s(x_2, b_2)) + r_\eta (x_3 - 2) \phi_\rho(x_2, b_2) \cdot (\phi_\eta^P(x_3, b_2) + \phi_\eta^T(x_3, b_2))] \alpha_s(t_f^2) h_f^2(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^2)] + [x_2 r_\rho \phi_\eta^A(x_3, b_2) (\phi_\rho^t(x_2, b_2) + \phi_\rho^s(x_2, b_2)) - x_3 r_\eta \phi_\rho(x_2, b_2) \cdot (\phi_\eta^P(x_3, b_2) + \phi_\eta^T(x_3, b_2))] \alpha_s(t_f^1) h_f^1(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^1)] \} . \quad (26)$$

$$\begin{aligned}
M_{a\eta}^{P_2} = & -\frac{32}{\sqrt{6}}\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_{B_s}(x_1, b_1) \\
& \cdot \left\{ [x_3 \phi_\rho(x_2, b_2) \phi_\eta^A(x_3, b_2) + r_\rho r_\eta \phi_\eta^P(x_3, b_2) ((x_2 + x_3 + 2) \phi_\rho^s(x_2, b_2) \right. \\
& - (x_2 - x_3) \phi_\rho^t(x_2, b_2)) + r_\rho r_\eta \phi_\eta^T(x_3, b_2) ((x_3 - x_2) \phi_\rho^s(x_2, b_2) + (x_2 \\
& + x_3 - 2) \phi_\rho^t(x_2, b_2))] \alpha_s(t_f^2) h_f^2(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^2)] \\
& + [(-x_2) \phi_\rho(x_2, b_2) \phi_\eta^A(x_3, b_2) + r_\rho r_\eta ((x_3 - x_2) (\phi_\eta^P(x_3, b_2) \phi_\rho^t(x_2, b_2) \\
& + \phi_\eta^T(x_3, b_2) \phi_\rho^s(x_2, b_2)) - (x_2 + x_3) (\phi_\eta^P(x_3, b_2) \phi_\rho^s(x_2, b_2) + \phi_\eta^T(x_3, b_2) \\
& \phi_\rho^t(x_2, b_2))] \alpha_s(t_f^1) h_f^1(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^1)] \} . \quad (27)
\end{aligned}$$

The factorizable annihilation diagrams 1(g) and 1(h) involve only ρ and η wave functions. Again decay amplitude $F_{a\eta}$ is for $(V - A)(V - A)$ type operators, while the $F_{a\eta}^{P_1}$ and $F_{a\eta}^{P_2}$ come from the $(V - A)(V + A)$ and $(S - P)(S + P)$ type operators, respectively

$$\begin{aligned}
F_{a\eta} = & 8\pi C_F f_{B_s} M_{B_s}^4 \int_0^1 dx_2 dx_3 \int_0^\infty b_2 db_2 b_3 db_3 \\
& \cdot \left\{ [x_3 \phi_\rho(x_2, b_2) \phi_\eta^A(x_3, b_3) + 2r_\rho r_\eta \phi_\rho^s(x_2, b_2) ((1 + x_3) \phi_\eta^P(x_3, b_3) \right. \\
& + (x_3 - 1) \phi_\eta^T(x_3, b_3))] \alpha_s(t_e^3) h_a(x_2, x_3, b_2, b_3) \exp[-S_{gh}(t_e^3)] \\
& - [x_2 \phi_\rho(x_2, b_2) \phi_\eta^A(x_3, b_3) + 2r_\rho r_\eta \phi_\eta^P(x_3, b_3) ((1 + x_2) \phi_\rho^s(x_2, b_2) \\
& + (x_2 - 1) \phi_\rho^t(x_2, b_2))] \alpha_s(t_e^4) h_a(x_3, x_2, b_3, b_2) \exp[-S_{gh}(t_e^4)] \} , \quad (28)
\end{aligned}$$

$$F_{a\eta}^{P_1} = -F_{a\eta}, \quad (29)$$

$$\begin{aligned}
F_{a\eta}^{P_2} = & -16\pi C_F f_{B_s} M_{B_s}^4 \int_0^1 dx_2 dx_3 \int_0^\infty b_2 db_2 b_3 db_3 \\
& \cdot \left\{ [2r_\rho \phi_\rho^s(x_2, b_2) \phi_\eta^A(x_3, b_3) + r_\eta x_3 \phi_\rho(x_2, b_2) (\phi_\eta^P(x_3, b_3) \right. \\
& - \phi_\eta^T(x_3, b_3))] \alpha_s(t_e^3) h_a(x_2, x_3, b_2, b_3) \exp[-S_{gh}(t_e^3)] \\
& + [x_2 r_\rho \phi_\eta^A(x_3, b_3) (\phi_\rho^s(x_2, b_2) - \phi_\rho^t(x_2, b_2)) + 2r_\eta \phi_\rho(x_2, b_2) \\
& \cdot \phi_\eta^P(x_3, b_3)] \alpha_s(t_e^4) h_a(x_3, x_2, b_3, b_2) \exp[-S_{gh}(t_e^4)] \} . \quad (30)
\end{aligned}$$

When we exchange the position of ρ and $\eta^{(\prime)}$ mesons in Fig. 1, then only 4 annihilation diagrams 1(e)-1(h) can contribute to $B \rightarrow \rho^0 \eta^{(\prime)}$ decays. The corresponding decay amplitudes from the non-factorizable annihilation diagrams 1(e) and 1(f), where it is the ρ^0 meson who picks up the spectator s quark, can be written as

$$\begin{aligned}
M_{a\rho} = & -\frac{32}{\sqrt{6}}\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_{B_s}(x_1, b_1) \\
& \cdot \left\{ [x_3 \phi_\rho(x_3, b_2) \phi_\eta^A(x_2, b_2) + r_\rho r_\eta ((x_3 - x_2) (\phi_\eta^P(x_2, b_2) \phi_\rho^t(x_3, b_2) \right. \\
& + \phi_\eta^T(x_2, b_2) \phi_\rho^s(x_3, b_2)) + (x_2 + x_3) (\phi_\eta^P(x_2, b_2) \phi_\rho^s(x_3, b_2) + \phi_\eta^T(x_2, b_2) \\
& \cdot \phi_\rho^t(x_3, b_2))] \alpha_s(t_f^1) h_f^1(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^1)] - [x_2 \phi_\rho(x_3, b_2) \\
& \cdot \phi_\eta^A(x_2, b_2) + r_\rho r_\eta ((x_2 - x_3) (\phi_\eta^P(x_2, b_2) \phi_\rho^t(x_3, b_2) + \phi_\eta^T(x_2, b_2) \\
& \cdot \phi_\rho^s(x_3, b_2))] + r_\rho r_\eta ((2 + x_2 + x_3) \phi_\eta^P(x_2, b_2) \phi_\rho^s(x_3, b_2) - (2 - x_2 - x_3) \\
& \cdot \phi_\eta^T(x_2, b_2) \phi_\rho^t(x_3, b_2))] \alpha_s(t_f^2) h_f^2(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^2)] \} , \quad (31)
\end{aligned}$$

$$\begin{aligned}
M_{a\rho}^{P_1} = & \frac{32}{\sqrt{6}}\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_{B_s}(x_1, b_1) \\
& \cdot \left\{ \left[x_2 r_\eta \phi_\rho(x_3, b_2) \left(\phi_\eta^P(x_2, b_2) + \phi_\eta^T(x_2, b_2) \right) - x_3 r_\rho \left(\phi_\rho^s(x_3, b_2) \right. \right. \right. \\
& \left. \left. \left. + \phi_\rho^t(x_3, b_2) \right) \phi_\eta^A(x_2, b_2) \right] \alpha_s(t_f^1) h_f^1(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^1)] \right. \\
& \left. + \left[(2-x_2) r_\eta \phi_\rho(x_3, b_2) \left(\phi_\eta^P(x_2, b_2) + \phi_\eta^T(x_2, b_2) \right) - (2-x_3) r_\rho \left(\phi_\rho^s(x_3, b_2) \right. \right. \right. \\
& \left. \left. \left. + \phi_\rho^t(x_3, b_2) \right) \phi_\eta^A(x_2, b_2) \right] \alpha_s(t_f^2) h_f^2(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^2)] \right\} , \quad (32)
\end{aligned}$$

$$\begin{aligned}
M_{a\rho}^{P_2} = & \frac{32}{\sqrt{6}}\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_{B_s}(x_1, b_1) \\
& \cdot \left\{ \left[x_2 \phi_\rho(x_3, b_2) \phi_\eta^A(x_2, b_2) + r_\rho r_\eta \left((x_2 + x_3) \left(\phi_\eta^P(x_2, b_2) \phi_\rho^s(x_3, b_2) \right. \right. \right. \right. \\
& \left. \left. \left. + \phi_\eta^T(x_2, b_2) \phi_\rho^t(x_3, b_2) \right) + (x_2 - x_3) \left(\phi_\eta^P(x_2, b_2) \phi_\rho^t(x_3, b_2) + \phi_\eta^T(x_2, b_2) \right. \right. \right. \\
& \left. \left. \left. \cdot \phi_\rho^s(x_3, b_2) \right) \right] \alpha_s(t_f^1) h_f^1(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^1)] - \left[x_3 \phi_\rho(x_3, b_2) \right. \right. \\
& \left. \left. \cdot \phi_\eta^A(x_2, b_2) + r_\rho r_\eta \left((x_3 - x_2) \left(\phi_\eta^P(x_2, b_2) \phi_\rho^t(x_3, b_2) + \phi_\eta^T(x_2, b_2) \phi_\rho^s(x_3, b_2) \right) \right) \right. \right. \\
& \left. \left. + r_\rho r_\eta \left((2 + x_2 + x_3) \phi_\eta^P(x_2, b_2) \phi_\rho^s(x_3, b_2) + (x_2 + x_3 - 2) \phi_\eta^T(x_2, b_2) \right. \right. \right. \\
& \left. \left. \left. \cdot \phi_\rho^t(x_3, b_2) \right) \right] \alpha_s(t_f^2) h_f^2(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^2)] \right\} . \quad (33)
\end{aligned}$$

For the factorizable annihilation diagrams 1(g) and 1(h) after the exchange of ρ^0 and $\eta^{(0)}$ mesons, the corresponding decay amplitudes can be obtained directly through the links with their counterparts $F_{a\eta}$, $F_{a\eta}$ and $F_{a\eta}^{P_2}$

$$F_{a\rho} = -F_{a\eta}, \quad F_{a\rho}^{P_1} = F_{a\eta}, \quad F_{a\rho}^{P_2} = F_{a\eta}^{P_2}. \quad (34)$$

Combining the contributions from different diagrams, the total decay amplitude for $B_s \rightarrow \rho^0 \eta$ decay can be written as:

$$\begin{aligned}
\mathcal{M}(\rho^0 \eta) = & F_{e\eta} \left[\xi_u \left(C_1 + \frac{1}{3} C_2 \right) - \xi_t \left(\frac{3}{2} C_7 + \frac{1}{2} C_8 + \frac{3}{2} C_9 + \frac{1}{2} C_{10} \right) \right] F_2(\phi) \\
& + M_{e\eta} \left[\xi_u C_2 - \xi_t \left(-\frac{3}{2} C_8 + \frac{3}{2} C_{10} \right) \right] F_2(\phi) \\
& + (M_{a\eta} + M_{a\rho}) \left[\xi_u C_2 - \xi_t \frac{3}{2} C_{10} \right] F_1(\phi) + (M_{a\eta}^{P_2} + M_{a\rho}^{P_2}) \left[-\xi_t \frac{3}{2} C_8 \right] F_1(\phi) \\
& + (F_{a\eta} + F_{a\rho}) \left[\xi_u \left(C_1 + \frac{1}{3} C_2 \right) \right. \\
& \left. - \xi_t \left(-\frac{3}{2} C_7 - \frac{1}{2} C_8 + \frac{3}{2} C_9 + \frac{1}{2} C_{10} \right) \right] F_1(\phi), \quad (35)
\end{aligned}$$

where $\xi_u = V_{ub}^* V_{ud}$, $\xi_t = V_{tb}^* V_{td}$, and $F_1(\phi) = \cos \phi/2$ and $F_2(\phi) = -\sin \phi/\sqrt{2}$ are the mixing factors. The Wilson coefficients C_i should be calculated at the appropriate scale t by using the formulas as given in the Appendices of Ref. [27].

Following the same steps, one can derive the total decay amplitude for $B_s \rightarrow \omega\eta$ decay:

$$\begin{aligned}
\mathcal{M}(\omega\eta) = & F_{e\eta} \left[\xi_u \left(C_1 + \frac{1}{3}C_2 \right) - \xi_t \left(2C_3 + \frac{2}{3}C_4 + 2C_5 \right. \right. \\
& \left. \left. + \frac{2}{3}C_6 + \frac{1}{2}C_7 + \frac{1}{6}C_8 + \frac{1}{2}C_9 + \frac{1}{6}C_{10} \right) \right] F_2(\phi) \\
& + M_{e\eta} \left[\xi_u C_2 - \xi_t \left(2C_4 - 2C_6 - \frac{1}{2}C_8 + \frac{1}{2}C_{10} \right) \right] F_2(\phi) \\
& + (M_{a\eta} + M_{a\omega}) \left[\xi_u C_2 - \xi_t \left(2C_4 + \frac{1}{2}C_{10} \right) \right] F_1(\phi) \\
& + (M_{a\eta}^{P_2} + M_{a\omega}^{P_2}) \left[-\xi_t \left(2C_6 + \frac{1}{2}C_8 \right) \right] F_1(\phi) + (F_{a\eta} + F_{a\omega}) \left[\xi_u \left(C_1 + \frac{1}{3}C_2 \right) \right. \\
& \left. - \xi_t \left(2C_3 + \frac{2}{3}C_4 - 2C_5 - \frac{2}{3}C_6 - \frac{1}{2}C_7 - \frac{1}{6}C_8 + \frac{1}{2}C_9 + \frac{1}{6}C_{10} \right) \right] F_1(\phi). \quad (36)
\end{aligned}$$

The individual decay amplitudes ($F_{e\eta}, M_{e\eta}, \dots$) in Eq. (36) can be obtained easily from those as given for the case of $B_s \rightarrow \rho^0\eta$ decay by the simple replacements

$$f_\rho \longrightarrow f_\omega, \quad f_\rho^T \longrightarrow f_\omega^T, \quad m_\rho \longrightarrow m_\omega \quad (37)$$

Note that the difference in the quark components of the two vector mesons ρ^0 and ω has been taken into account.

For $B_s \rightarrow \phi\eta^{(\prime)}$ decays, all eight Feynman diagrams when the $\eta^{(\prime)}$ or ϕ meson picks up the spectator s quark. For the later case, we firstly consider the factorizable diagrams 1(a) and 1(b). The decay amplitude $F_{e\phi}$ induced by inserting the $(V-A)(V+A)$ operators is

$$\begin{aligned}
F_{e\phi} = & -8\pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \phi_{B_s}(x_1, b_1) \\
& \cdot \left\{ \left[(1+x_3)\phi_\phi(x_3, b_3) + r_\phi(1-2x_3)(\phi_\phi^s(x_3, b_3) + \phi_\phi^t(x_3, b_3)) \right] \right. \\
& \alpha_s(t_e^1) h_e(x_1, x_3, b_1, b_3) \exp[-S_{ab}(t_e^1)] \\
& \left. + 2r_\phi \phi_\phi^s(x_3, b_3) \alpha_s(t_e^2) h_e(x_3, x_1, b_3, b_1) \exp[-S_{ab}(t_e^2)] \right\}, \quad (38)
\end{aligned}$$

where $r_\phi = m_\phi/M_{B_s}$. One can also extract out the $B_s \rightarrow \phi$ form factor $A_0^{B_s \rightarrow \phi}$ from the diagrams 1(a) and 1(b) when ϕ meson picks up the spectator s quark. For other diagrams, the relevant decay amplitudes can be written as

$$F_{e\phi}^{P_1} = -F_{e\phi}, \quad (39)$$

$$\begin{aligned}
F_{e\phi}^{P_2} = & -16\pi C_F M_{B_s}^4 r_\eta \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \phi_{B_s}(x_1, b_1) \\
& \cdot \left\{ \left[\phi_\phi(x_3, b_3) + r_\phi((x_3+2)\phi_\phi^s(x_3, b_3) - x_3\phi_\phi^t(x_3, b_3)) \right] \right. \\
& \cdot \alpha_s(t_e^1) h_e(x_1, x_3, b_1, b_3) \exp[-S_{ab}(t_e^1)] + (x_1\phi_\phi(x_3, b_3) \\
& \left. + 2r_\phi \phi_\phi^s(x_3, b_3)) \alpha_s(t_e^2) h_e(x_3, x_1, b_3, b_1) \exp[-S_{ab}(t_e^2)] \right\}, \quad (40)
\end{aligned}$$

$$\begin{aligned}
M_{e\phi} = & \frac{32}{\sqrt{6}} \pi C_F M_{B_s}^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \\
& \cdot \phi_{B_s}(x_1, b_1) \phi_\eta^A(x_2, b_2) \left\{ x_3 \left[\phi_\phi(x_3, b_1) - 2r_\phi \phi_\phi^t(x_3, b_1) \right] \right. \\
& \cdot \alpha_s(t_f) h_f(x_1, x_2, x_3, b_1, b_2) \exp[-S_{cd}(t_f)] \left. \right\} ,
\end{aligned} \tag{41}$$

$$M_{e\phi}^{P_1} = 0, \quad M_{e\phi}^{P_2} = M_{e\phi}. \tag{42}$$

Finally, the total decay amplitude of $B_s \rightarrow \phi\eta$ decay can be written as

$$\begin{aligned}
\mathcal{M}(\phi\eta) = & \sqrt{2} F_{e\phi} \left\{ \left[\xi_u \left(C_1 + \frac{1}{3} C_2 \right) \right. \right. \\
& - \xi_t \left(2C_3 + \frac{2}{3} C_4 - 2C_5 - \frac{2}{3} C_6 - \frac{1}{2} C_7 - \frac{1}{6} C_8 + \frac{1}{2} C_9 + \frac{1}{6} C_{10} \right) \left. \right] f_q F_1(\phi) \\
& - \xi_t \left(\frac{4}{3} C_3 + \frac{4}{3} C_4 - C_5 - \frac{1}{3} C_6 + \frac{1}{2} C_7 + \frac{1}{6} C_8 - \frac{2}{3} C_9 - \frac{2}{3} C_{10} \right) f_s F_2(\phi) \left. \right\} \\
& + \sqrt{2} F_{e\phi}^{P_2} \left[-\xi_t \left(\frac{1}{3} C_5 + C_6 - \frac{1}{6} C_7 - \frac{1}{2} C_8 \right) \right] f_s F_2(\phi) \\
& + \sqrt{2} F_{e\eta} \left[-\xi_t \left(\frac{4}{3} C_3 + \frac{4}{3} C_4 + C_5 + \frac{1}{3} C_6 - \frac{1}{2} C_7 - \frac{1}{6} C_8 - \frac{2}{3} C_9 - \frac{2}{3} C_{10} \right) \right] F_2(\phi) \\
& + \sqrt{2} M_{e\phi} \left\{ \left[\xi_u (C_2) - \xi_t \left(2C_4 + 2C_6 + \frac{1}{2} C_8 + \frac{1}{2} C_{10} \right) \right] F_1(\phi) \right. \\
& - \xi_t \left(C_3 + C_4 + C_6 - \frac{1}{2} C_8 - \frac{1}{2} C_9 - \frac{1}{2} C_{10} \right) F_2(\phi) \left. \right\} \\
& + \sqrt{2} M_{e\eta} \left[-\xi_t \left(C_3 + C_4 - C_6 + \frac{1}{2} C_8 - \frac{1}{2} C_9 - \frac{1}{2} C_{10} \right) \right] F_2(\phi) \\
& + \sqrt{2} M_{e\eta}^{P_1} \left[-\xi_t \left(C_5 - \frac{1}{2} C_7 \right) \right] F_2(\phi) \\
& + \sqrt{2} (M_{a\eta} + M_{a\phi}) \left[-\xi_t \left(C_3 + C_4 - \frac{1}{2} C_9 - \frac{1}{2} C_{10} \right) \right] F_2(\phi) \\
& + \sqrt{2} (M_{a\eta}^{P_1} + M_{a\phi}^{P_1}) \left[-\xi_t \left(C_5 - \frac{1}{2} C_7 \right) \right] F_2(\phi) \\
& + \sqrt{2} (M_{a\eta}^{P_2} + M_{a\phi}^{P_2}) \left[-\xi_t \left(C_6 - \frac{1}{2} C_8 \right) \right] F_2(\phi) \\
& + \sqrt{2} (F_{a\eta} + F_{a\phi}) \left[-\xi_t \left(\frac{4}{3} C_3 + \frac{4}{3} C_4 - C_5 \right. \right. \\
& \left. \left. - \frac{1}{3} C_6 + \frac{1}{2} C_7 + \frac{1}{6} C_8 - \frac{2}{3} C_9 - \frac{2}{3} C_{10} \right) \right] F_2(\phi) \\
& + \sqrt{2} (F_{a\eta}^{P_2} + F_{a\phi}^{P_2}) \left[-\xi_t \left(\frac{1}{3} C_5 + C_6 - \frac{1}{6} C_7 - \frac{1}{2} C_8 \right) \right] F_2(\phi).
\end{aligned} \tag{43}$$

The individual decay amplitudes $(F_{e\eta}, M_{e\eta}, \dots)$ in Eq. (43) can be obtained easily from

those as given for the case of $B_s \rightarrow \rho^0 \eta$ decay by the simple replacements

$$\begin{aligned} f_\rho &\longrightarrow f_\phi, & f_\rho^T &\longrightarrow f_\phi^T, & m_\rho &\longrightarrow m_\phi \\ \phi_\rho &\longrightarrow \phi_\phi, & \phi_\rho^t &\longrightarrow \phi_\phi^t, & \phi_\rho^s &\longrightarrow \phi_\phi^s \end{aligned} \quad (44)$$

The total decay amplitudes for $B_s \rightarrow \rho^0 \eta'$, $B_s \rightarrow \omega \eta'$ and $B_s \rightarrow \phi \eta'$ can be obtained easily from Eqs.(35), (36) and (43) by the following simple replacements:

$$\begin{aligned} F_1(\phi) = \cos \phi &\longrightarrow F'_1(\phi) = \sin \phi, \\ F_2(\phi) = -\sin \phi / \sqrt{2} &\longrightarrow F'_2(\phi) = \cos \phi / \sqrt{2}. \end{aligned} \quad (45)$$

Note that the possible gluonic component of η' meson has been neglected here. We will estimate its effects in next section.

IV. NUMERICAL RESULTS AND DISCUSSIONS

A. Input parameters and wave functions

We use the following input parameters in the numerical calculations

$$\begin{aligned} \Lambda_{\overline{\text{MS}}}^{(f=4)} &= 250\text{MeV}, & f_\rho &= 205\text{MeV}, & f_\rho^T &= 160\text{MeV}, \\ m_0^{\eta_{d\bar{d}}} &= 1.4\text{GeV}, & f_{B_s} &= 230\text{MeV}, & f_\pi &= 130\text{MeV}, \\ m_\omega &= 0.782\text{GeV}, & f_\omega &= 195\text{MeV}, & f_\omega^T &= 140\text{MeV}, \\ m_\phi &= 1.02\text{GeV}, & f_\phi &= 237\text{MeV}, & f_\phi^T &= 220\text{MeV}, \\ m_\rho &= 0.770\text{GeV}, & M_{B_s} &= 5.37\text{GeV}, & M_W &= 80.42\text{GeV}. \end{aligned} \quad (46)$$

For the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements, here we adopt the Wolfenstein parametrization for the CKM matrix up to $\mathcal{O}(\lambda^5)$ with the parameters $\lambda = 0.2272$, $A = 0.818$, $\bar{\rho} = 0.221$ and $\bar{\eta} = 0.340$ [19].

B_s meson is different from B meson due to the heavier strange quark (compare to u, d quark) which induces the SU(3) symmetry-breaking effect. This effect is considered to be small and the distribution amplitude of B_s meson (given in the following formula) should be similar to that of the B meson,

$$\phi_{B_s}(x, b) = N_{B_s} x^2 (1-x)^2 \exp \left[-\frac{M_{B_s}^2 x^2}{2\omega_{B_s}^2} - \frac{1}{2}(\omega_{B_s} b)^2 \right], \quad (47)$$

where ω_{B_s} is a free parameter in nature. After considering the constraints from some B_s non-leptonic decays [28], we here use $\omega_{B_s} = 0.55 \pm 0.05$ in the numerical calculation.

For the light meson wave function, we neglect the b dependant part, which is not important in numerical analysis. We choose the wave function of $\rho(\omega)$ meson similar to

the pion case [29]:

$$\phi_{\rho(\omega)}(x) = \frac{3}{\sqrt{6}} f_{\rho(\omega)} x(1-x) \left[1 + 0.18 C_2^{\frac{3}{2}}(2x-1) \right], \quad (48)$$

$$\begin{aligned} \phi_{\rho(\omega)}^t(x) = & \frac{f_{\rho(\omega)}^T}{2\sqrt{6}} \left\{ 3(2x-1)^2 + 0.3(2x-1)^2 [5(2x-1)^2 - 3] \right. \\ & \left. + 0.21[3 - 30(2x-1)^2 + 35(2x-1)^4] \right\}, \end{aligned} \quad (49)$$

$$\phi_{\rho(\omega)}^s(x) = \frac{3}{2\sqrt{6}} f_{\rho(\omega)}^T (1-2x) \left[1 + 0.76(10x^2 - 10x + 1) \right]. \quad (50)$$

For the wave function of ϕ meson, we use [29]:

$$\phi_\phi(x) = \frac{3}{\sqrt{6}} f_\phi x(1-x), \quad (51)$$

$$\phi_\phi^t(x) = \frac{f_\phi^T}{2\sqrt{6}} \left\{ 3(1-2x)^2 + 1.68 C_4^{\frac{1}{2}}(1-2x) + 0.69 \left[1 + (1-2x) \ln \frac{x}{1-x} \right] \right\}, \quad (52)$$

$$\phi_\phi^s(x) = \frac{f_\phi^T}{4\sqrt{6}} \left[3(1-2x)(4.5 - 11.2x + 11.2x^2) + 1.38 \ln \frac{x}{1-x} \right], \quad (53)$$

where the Gegenbauer polynomials are defined by

$$C_2^{\frac{3}{2}}(\xi) = \frac{3}{2} (5\xi^2 - 1), \quad (54)$$

$$C_4^{\frac{1}{2}}(\xi) = \frac{1}{8} (35\xi^4 - 30\xi^2 + 3). \quad (55)$$

For η meson's wave function, $\phi_{\eta d\bar{d}}^A$, $\phi_{\eta d\bar{d}}^P$ and $\phi_{\eta d\bar{d}}^T$ represent the axial vector, pseudoscalar and tensor components of the wave function respectively, for which we utilize the results from the light-cone sum rule [30] including twist-3 contribution:

$$\begin{aligned} \phi_{\eta d\bar{d}}^A(x) = & \frac{3}{\sqrt{2N_c}} f_q x(1-x) \left\{ 1 + a_2^{\eta d\bar{d}} \frac{3}{2} [5(1-2x)^2 - 1] \right. \\ & \left. + a_4^{\eta d\bar{d}} \frac{15}{8} [21(1-2x)^4 - 14(1-2x)^2 + 1] \right\}, \\ \phi_{\eta d\bar{d}}^P(x) = & \frac{1}{2\sqrt{2N_c}} f_q \left\{ 1 + \frac{1}{2} \left(30\eta_3 - \frac{5}{2} \rho_{\eta d\bar{d}}^2 \right) [3(1-2x)^2 - 1] \right. \\ & \left. + \frac{1}{8} \left(-3\eta_3\omega_3 - \frac{27}{20} \rho_{\eta d\bar{d}}^2 - \frac{81}{10} \rho_{\eta d\bar{d}}^2 a_2^{\eta d\bar{d}} \right) [35(1-2x)^4 - 30(1-2x)^2 + 3] \right\}, \\ \phi_{\eta d\bar{d}}^T(x) = & \frac{3}{\sqrt{2N_c}} f_q (1-2x) \\ & \cdot \left[\frac{1}{6} + (5\eta_3 - \frac{1}{2} \eta_3 \omega_3 - \frac{7}{20} \rho_{\eta d\bar{d}}^2 - \frac{3}{5} \rho_{\eta d\bar{d}}^2 a_2^{\eta d\bar{d}}) (10x^2 - 10x + 1) \right], \end{aligned} \quad (56)$$

with the updated Gegenbauer moments[31]

$$\begin{aligned} a_2^{\eta d\bar{d}} &= 0.115, \quad a_4^{\eta d\bar{d}} = -0.015, \\ \rho_{\eta d\bar{d}} &= m_\pi/m_0^{\eta d\bar{d}}, \quad \eta_3 = 0.015, \quad \omega_3 = -3.0 \end{aligned} \quad (57)$$

We assume that the wave function of $u\bar{u}$ is same as the wave function of $d\bar{d}$. For the wave function of the $s\bar{s}$ components, we also use the same form as $d\bar{d}$ but with $m_0^{\eta_{s\bar{s}}}$ and f_s instead of $m_0^{\eta_{d\bar{d}}}$ and f_q , respectively.

B. Branching ratios

For the decays we have considered here, the decay amplitudes in Eqs. (35) , (36) and (43) can be rewritten as

$$\mathcal{M} = V_{ub}^* V_{us} T - V_{tb}^* V_{ts} P = V_{ub}^* V_{us} T [1 + z e^{i(\gamma+\delta)}], \quad (58)$$

where

$$z = \left| \frac{V_{tb}^* V_{ts}}{V_{ub}^* V_{us}} \right| \left| \frac{P}{T} \right| \quad (59)$$

is the ratio of penguin to tree contributions, $\gamma = \arg \left[-\frac{V_{ts} V_{tb}^*}{V_{us} V_{ub}^*} \right]$ is the weak phase (one of the three CKM angles), and δ is the relative strong phase between tree (T) and penguin (P) diagrams.

From Eq. (58), it is easy to write the decay amplitude for the corresponding charge conjugated decay mode

$$\overline{\mathcal{M}} = V_{ub} V_{us}^* T - V_{tb} V_{ts}^* P = V_{ub} V_{us}^* T [1 + z e^{i(-\gamma+\delta)}]. \quad (60)$$

Therefore the CP-averaged branching ratio is

$$Br = (|\mathcal{M}|^2 + |\overline{\mathcal{M}}|^2)/2 = |V_{ub} V_{us}^* T|^2 [1 + 2z \cos \gamma \cos \delta + z^2], \quad (61)$$

where the ratio z and the strong phase δ have been defined in Eqs.(58) and (59).

Using the wave functions and the input parameters as specified in previous sections, it is straightforward to calculate the CP averaged branching ratios for the considered decays:

$$\begin{aligned} Br(B_s \rightarrow \rho^0 \eta) &= [0.07_{-0.02}^{+0.03}(\omega_{B_s})_{-0.03}^{+0.05}(m_s)_{-0.00}^{+0.00}(\gamma)] \times 10^{-6}, \\ Br(B_s \rightarrow \rho^0 \eta') &= [0.10_{-0.03}^{+0.04}(\omega_{B_s})_{-0.04}^{+0.07}(m_s)_{-0.01}^{+0.00}(\gamma)] \times 10^{-6}, \end{aligned} \quad (62)$$

$$\begin{aligned} Br(B_s \rightarrow \omega \eta) &= [0.21_{-0.03}^{+0.04}(\omega_{B_s})_{+0.00}^{+0.14}(m_s) \pm 0.00(\gamma)] \times 10^{-7}, \\ Br(B_s \rightarrow \omega \eta') &= [0.13_{-0.03}^{+0.04}(\omega_{B_s})_{-0.03}^{+0.05}(m_s)_{-0.01}^{+0.00}(\gamma)] \times 10^{-6}, \end{aligned} \quad (63)$$

$$\begin{aligned} Br(B_s \rightarrow \phi \eta) &= [2.66_{-0.74}^{+1.10}(\omega_{B_s})_{-0.78}^{+1.45}(m_s)_{-0.00}^{+0.01}(\gamma)] \times 10^{-5}, \\ Br(B_s \rightarrow \phi \eta') &= [2.00_{-0.54}^{+0.80}(\omega_{B_s})_{-0.73}^{+1.42}(m_s)_{-0.00}^{+0.00}(\gamma)] \times 10^{-5}, \end{aligned} \quad (64)$$

where the main errors are induced by the uncertainty of $\omega_{B_s} = 0.55 \pm 0.05$ GeV, $m_s = 130 \pm 30$ MeV and $\gamma = 60^\circ \pm 20^\circ$, respectively.

As a comparison, we also list here the theoretical predictions based on the QCD factorization approach as given in Ref. [7]:

$$\begin{aligned} Br(B_s \rightarrow \rho^0 \eta) &= [0.122 - 0.151] \times 10^{-6}, \\ Br(B_s \rightarrow \rho^0 \eta') &= [0.123 - 0.160] \times 10^{-6}, \end{aligned} \quad (65)$$

$$\begin{aligned} Br(B_s \rightarrow \omega \eta) &= [0.006 - 0.025] \times 10^{-6}, \\ Br(B_s \rightarrow \omega \eta') &= [0.010 - 0.075] \times 10^{-6}, \end{aligned} \quad (66)$$

$$\begin{aligned} Br(B_s \rightarrow \phi \eta) &= [0.088 - 0.417] \times 10^{-6}, \\ Br(B_s \rightarrow \phi \eta') &= [0.024 - 0.149] \times 10^{-6}. \end{aligned} \quad (67)$$




FIG. 2: The CP averaged branching ratio (in unit of 10^{-6}) of $B_s \rightarrow \rho^0 \eta$ decay as a function of CKM angle γ . (a) is for $m_s = 130$ MeV and $\omega_{B_s} = 0.50$ GeV (dotted curve), 0.55 GeV (solid curve) and 0.60 GeV (dashed curve); and (b) for $\omega_{B_s} = 0.55$ GeV, and $m_s = 100$ MeV (dotted curve), 130 MeV (solid curve) and 160 MeV (dashed curve).




FIG. 3: The same as Fig. 2 but for $B_s \rightarrow \rho^0 \eta'$ decay.

One can see that the pQCD predictions are basically consistent with the corresponding QCDF predictions for the first four decays, but much larger than the QCDF predictions for $B_s \rightarrow \phi \eta$ and $\phi \eta'$ decays. The difference is about two orders! Further study about this difference is under way.

In Figs. 2-7, we present the PQCD predictions of the CP averaged branching ratios of the considered decays and their dependence on the variations of ω_{B_s} , m_s and the CKM angle γ .




FIG. 4: The same as Fig. 2 but for $B_s \rightarrow \omega\eta$ decay.




FIG. 5: The same as Fig. 2 but for $B_s \rightarrow \omega\eta'$ decay.

C. CP-violating asymmetries

Now we turn to the evaluation of the CP-violating asymmetries of the considered decays in PQCD approach. For these neutral decay modes, the effects of $B_s^0 - \bar{B}_s^0$ mixing should be considered.

For B_s^0 meson decays, the CP-violating asymmetry of $B_s^0(\bar{B}_s^0) \rightarrow f_{CP}$ decay is time dependent and can be defined as

$$\begin{aligned}
 A_{CP} &\equiv \frac{\Gamma(\bar{B}_s^0(\Delta t) \rightarrow f_{CP}) - \Gamma(B_s^0(\Delta t) \rightarrow f_{CP})}{\Gamma(\bar{B}_s^0(\Delta t) \rightarrow f_{CP}) + \Gamma(B_s^0(\Delta t) \rightarrow f_{CP})} \\
 &= A_{CP}^{dir} \cos(\Delta m_s \Delta t) + A_{CP}^{mix} \sin(\Delta m_s \Delta t),
 \end{aligned} \tag{68}$$

where Δm_s is the mass difference between the two B_s^0 mass eigenstates, $\Delta t = t_{CP} - t_{tag}$ is the time difference between the tagged B_s^0 ($\bar{B}_s^0$) and the accompanying $\bar{B}_s^0$ (B_s^0) with opposite b flavor decaying to the final CP-eigenstate f_{CP} at the time t_{CP} . The direct and




FIG. 6: The same as Fig. 2 but for $B_s \rightarrow \phi\eta$ decay.




FIG. 7: The same as Fig. 2 but for $B_s \rightarrow \phi\eta'$ decay.

mixing induced CP-violating asymmetries A_{CP}^{dir} and A_{CP}^{mix} can be written as

$$A_{CP}^{dir} = \frac{|\lambda_{CP}|^2 - 1}{1 + |\lambda_{CP}|^2}, \quad A_{CP}^{mix} = \frac{2Im(\lambda_{CP})}{1 + |\lambda_{CP}|^2}, \quad (69)$$

where the CP-violating parameter λ_{CP} is

$$\lambda_{CP} = \frac{V_{tb}^* V_{ts} \langle \bar{f}_{CP} | H_{eff} | \bar{B}_s^0 \rangle}{V_{tb} V_{ts}^* \langle f_{CP} | H_{eff} | B_s^0 \rangle} = e^{2i\gamma} \frac{1 + ze^{i(\delta-\gamma)}}{1 + ze^{i(\delta+\gamma)}}. \quad (70)$$

Here the ratio z and the strong phase δ have been defined previously. In PQCD approach, since both z and δ are calculable, it is easy to find the numerical values of A_{CP}^{dir} and A_{CP}^{mix} for the considered decay processes.

The PQCD predictions for the direct and mixing-induced CP-violating asymmetries

are:

$$\begin{aligned} A_{CP}^{dir}(B_s \rightarrow \rho^0 \eta) &= [4.9_{-0.1}^{+0.4}(\omega_{B_s})_{-0.5}^{+0.4}(m_s)_{-1.2}^{+0.6}(\gamma)] \times 10^{-2}, \\ A_{CP}^{dir}(B_s \rightarrow \rho^0 \eta') &= [-25.3_{-2.0}^{+1.8}(\omega_{B_s})_{-8.7}^{+7.7}(m_s)_{-5.3}^{+7.4}(\gamma)] \times 10^{-2}, \end{aligned} \quad (71)$$

$$\begin{aligned} A_{CP}^{dir}(B_s \rightarrow \omega \eta) &= [7.6_{-5.4}^{+5.1}(\omega_{B_s})_{-12.8}^{+11.1}(m_s)_{-2.1}^{+1.3}(\gamma)] \times 10^{-2}, \\ A_{CP}^{dir}(B_s \rightarrow \omega \eta') &= [-7.6_{-0.2}^{+0.4}(\omega_{B_s})_{-2.8}^{+3.1}(m_s)_{-1.9}^{+2.3}(\gamma)] \times 10^{-2}. \end{aligned} \quad (72)$$

$$\begin{aligned} A_{CP}^{dir}(B_s \rightarrow \phi \eta) &= [0.5_{-0.1}^{+0.0}(\omega_{B_s})_{-0.1}^{+0.1}(m_s)_{-0.1}^{+0.1}(\gamma)] \times 10^{-2}, \\ A_{CP}^{dir}(B_s \rightarrow \phi \eta') &= [-0.5_{-0.0}^{+0.1}(\omega_{B_s})_{-0.1}^{+0.1}(m_s)_{-0.1}^{+0.1}(\gamma)] \times 10^{-2}, \end{aligned} \quad (73)$$

$$\begin{aligned} A_{CP}^{mix}(B_s \rightarrow \rho^0 \eta) &= [-4.2_{-4.2}^{+3.8}(\omega_{B_s})_{-13.1}^{+10.8}(m_s)_{-0.6}^{+1.1}(\gamma)] \times 10^{-2}, \\ A_{CP}^{mix}(B_s \rightarrow \rho^0 \eta') &= [18.7_{+0.5}^{+0.3}(\omega_{B_s})_{-3.9}^{+3.7}(m_s)_{-4.4}^{+1.6}(\gamma)] \times 10^{-2}, \end{aligned} \quad (74)$$

$$\begin{aligned} A_{CP}^{mix}(B_s \rightarrow \omega \eta) &= [8.0_{+0.7}^{+0.7}(\omega_{B_s})_{+5.5}^{+6.4}(m_s)_{-2.1}^{+1.1}(\gamma)] \times 10^{-2}, \\ A_{CP}^{mix}(B_s \rightarrow \omega \eta') &= [24.6_{-0.6}^{+1.1}(\omega_{B_s})_{+0.4}^{+0.6}(m_s)_{-6.8}^{+4.4}(\gamma)] \times 10^{-2}. \end{aligned} \quad (75)$$

$$\begin{aligned} A_{CP}^{mix}(B_s \rightarrow \phi \eta) &= [-0.5_{-0.0}^{+0.0}(\omega_{B_s})_{-0.1}^{+0.1}(m_s)_{-0.1}^{+0.1}(\gamma)] \times 10^{-2}, \\ A_{CP}^{mix}(B_s \rightarrow \phi \eta') &= [0.5_{-0.0}^{+0.1}(\omega_{B_s})_{-0.1}^{+0.1}(m_s)_{-0.1}^{+0.1}(\gamma)] \times 10^{-2}, \end{aligned} \quad (76)$$

where the dominant errors come from the variation of $\omega_{B_s} = 0.55 \pm 0.05$, $m_s = 130 \pm 30$ Mev, $\gamma = 60^\circ \pm 20^\circ$. One can see from these numerical results that the CP-violating asymmetries of the considered decay modes are generally not large in size. This is rather different with the cases of B_d and B_u meson decays, where the CP-violating asymmetries are generally large in magnitude.

As a simple comparison, we show the QCDF predictions for the direct CP-violating asymmetries of the six considered decays as given in Ref. [7]:

$$\begin{aligned} A_{CP}^{dir}(B_s \rightarrow \rho^0 \eta) &= -16.9 \times 10^{-2} \\ A_{CP}^{dir}(B_s \rightarrow \rho^0 \eta') &= -33.0 \times 10^{-2}, \end{aligned} \quad (77)$$

$$\begin{aligned} A_{CP}^{dir}(B_s \rightarrow \omega \eta) &= 0.76 \times 10^{-2}, \\ A_{CP}^{dir}(B_s \rightarrow \omega \eta') &= 4.4 \times 10^{-2}, \end{aligned} \quad (78)$$

$$\begin{aligned} A_{CP}^{dir}(B_s \rightarrow \phi \eta) &= 23.2 \times 10^{-2}, \\ A_{CP}^{dir}(B_s \rightarrow \phi \eta') &= -58.3 \times 10^{-2}. \end{aligned} \quad (79)$$

Since the relevant measurements are not available at present, and the theoretical uncertainties in both pQCD and QCDF factorization approaches are still very large, it is too early to make a meaningful comparison now. One has to wait for the starting of experimental measurements and improvements of the theoretical calculations.

In Fig. 6 we show the γ -dependence of the direct and mixing induced CP-violating asymmetries for $B_s^0 \rightarrow \rho^0 \eta^{(\prime)}$ and $\omega \eta^{(\prime)}$ decays. Since the pQCD predictions of the CP asymmetries are sensitive to many input parameters, the lines in Fig. 6 should be broadened accordingly.

D. Effects of possible gluonic component of η'

Up to now, we have not considered the possible contributions to the branching ratios and CP-violating asymmetries of $B \rightarrow (\rho^0, \omega, \phi) \eta^{(\prime)}$ decays induced by the possible gluonic component of $\eta^{(\prime)}$ [18, 32].




FIG. 8: The direct and mixing-induced CP asymmetry (in percentage) of $B_s^0 \rightarrow \rho^0 \eta$ (solid curve), $\rho^0 \eta'$ (dotted curve), $\omega \eta$ (dashed-curve), and $\omega \eta'$ (dash-dot-dot curve) decay as a function of CKM angle γ .

By using the formulae as given in Ref. [22], we calculate the gluonic contributions to $B \rightarrow \eta$ and $B \rightarrow \eta'$ form factors. For $B \rightarrow (\rho, \omega, \phi) \eta$ decays, the gluonic contributions are negligibly small, less than 3%. For $B \rightarrow (\rho, \omega, \phi) \eta'$ decays, the gluonic contributions to the branching ratios are only about 10%. The central values of the pQCD predictions for $B \rightarrow (\rho, \omega, \phi) \eta'$ decays after the inclusion of gluonic contributions are the following

$$Br(B_s \rightarrow \rho^0 \eta') = [0.12_{-0.03}^{+0.04}(\omega_{B_s})_{-0.04}^{+0.07}(m_s)_{-0.01}^{+0.00}(\gamma)] \times 10^{-6}, \quad (80)$$

$$Br(B_s \rightarrow \omega \eta') = [0.14_{-0.03}^{+0.05}(\omega_{B_s})_{-0.03}^{+0.06}(m_s)_{-0.01}^{+0.01}(\gamma)] \times 10^{-6}, \quad (81)$$

$$Br(B_s \rightarrow \phi \eta') = [2.11_{-0.57}^{+0.84}(\omega_{B_s})_{-0.75}^{+1.45}(m_s)_{-0.00}^{+0.00}(\gamma)] \times 10^{-5}. \quad (82)$$

Again, the gluonic contributions are small in size, which is consistent with previous results for other B meson decays involving η or η' as one or two final state mesons [23, 24, 26, 33].

V. SUMMARY

In this paper, we calculated the branching ratios and CP-violating asymmetries of $B_s \rightarrow \rho^0 \eta^{(\prime)}$, $B_s \rightarrow \omega \eta^{(\prime)}$, and $B_s \rightarrow \phi \eta^{(\prime)}$ decays in the pQCD factorization approach.

Besides the usual factorizable diagrams, the non-factorizable and annihilation diagrams as shown in Fig. (1) are also calculated analytically. Although the non-factorizable and annihilation contributions are sub-leading for the branching ratios of the considered decays, but they are not negligible. Furthermore these diagrams provide the necessary strong phase required for a non-zero CP-violating asymmetry for the considered decays.

From our calculations and phenomenological analysis, we found the following results:

- For the branching ratios of the six considered decay modes, the pQCD predictions

are:

$$\begin{aligned} Br(B_s \rightarrow \rho^0 \eta) &= [0.07_{-0.04}^{+0.06}] \times 10^{-6}, \\ Br(B_s \rightarrow \rho^0 \eta') &= [0.10_{-0.05}^{+0.08}] \times 10^{-6}, \end{aligned} \quad (83)$$

$$\begin{aligned} Br(B_s \rightarrow \omega \eta) &= [0.02_{-0.003}^{+0.015}] \times 10^{-6}, \\ Br(B_s \rightarrow \omega \eta') &= [0.13_{-0.04}^{+0.06}] \times 10^{-6}, \end{aligned} \quad (84)$$

$$\begin{aligned} Br(B_s \rightarrow \phi \eta) &= [2.66_{-1.08}^{+1.82}] \times 10^{-5}, \\ Br(B_s \rightarrow \phi \eta') &= [2.00_{-0.91}^{+1.63}] \times 10^{-5}. \end{aligned} \quad (85)$$

Here the various errors as specified previously have been added in quadrature. The pQCD predictions for the first four decay channels agree well with those obtained by employing the QCDF approach. For last two decays, however, there are large differences between the pQCD and QCDF approach, which will be tested in the forthcoming LHC-b experiments.

- The gluonic contributions are small in size: less than 3% for $B \rightarrow (\rho, \omega, \phi)\eta$ decays, and about 10% for $B \rightarrow (\rho, \omega, \phi)\eta'$ decays.
- For the CP-violating asymmetries, the pQCD predictions are generally not large in size, but still have large theoretical uncertainties.

Acknowledgments

We are very grateful to Cai-Dian Lü, Hui-sheng Wang, Xin Liu and Ying Li for helpful discussions. This work is partly supported by the National Natural Science Foundation of China under Grant No.10575052, by the Specialized Research Fund for the doctoral Program of higher education (SRFDP) under Grant No. 20050319008 and by the Research Fund from Nanjing Normal University under Grant No. 214080A916.

APPENDIX A: RELATED FUNCTIONS

We show here the function h_i 's, coming from the Fourier transformations of $H^{(0)}$,

$$\begin{aligned} h_e(x_1, x_3, b_1, b_3) &= K_0(\sqrt{x_1 x_3} m_{B_s} b_1) [\theta(b_1 - b_3) K_0(\sqrt{x_3} m_{B_s} b_1) I_0(\sqrt{x_3} m_{B_s} b_3) \\ &\quad + \theta(b_3 - b_1) K_0(\sqrt{x_3} m_{B_s} b_3) I_0(\sqrt{x_3} m_{B_s} b_1)] S_t(x_3), \end{aligned} \quad (A1)$$

$$\begin{aligned} h_a(x_2, x_3, b_2, b_3) &= K_0(i\sqrt{x_2 x_3} m_{B_s} b_2) [\theta(b_3 - b_2) K_0(i\sqrt{x_3} m_{B_s} b_3) I_0(i\sqrt{x_3} m_{B_s} b_2) \\ &\quad + \theta(b_2 - b_3) K_0(i\sqrt{x_3} m_{B_s} b_2) I_0(i\sqrt{x_3} m_{B_s} b_3)] S_t(x_3), \end{aligned} \quad (A2)$$

$$\begin{aligned} h_f(x_1, x_2, x_3, b_1, b_2) &= \left\{ \theta(b_2 - b_1) I_0(M_{B_s} \sqrt{x_1 x_3} b_1) K_0(M_{B_s} \sqrt{x_1 x_3} b_2) \right. \\ &\quad \left. + (b_1 \leftrightarrow b_2) \right\} \cdot \left(\begin{array}{ll} K_0(M_{B_s} F_{(1)} b_2), & \text{for } F_{(1)}^2 > 0 \\ \frac{\pi i}{2} H_0^{(1)}(M_{B_s} \sqrt{|F_{(1)}^2|} b_2), & \text{for } F_{(1)}^2 < 0 \end{array} \right), \end{aligned} \quad (A3)$$

$$h_f^1(x_1, x_2, x_3, b_1, b_2) = \left\{ \theta(b_1 - b_2) K_0(i\sqrt{x_2 x_3} b_1 M_{B_s}) I_0(i\sqrt{x_2 x_3} b_2 M_{B_s}) \right. \\ \left. + (b_1 \leftrightarrow b_2) \right\} \cdot \left(\begin{array}{ll} K_0(M_{B_s} F_{(2)} b_1), & \text{for } F_{(2)}^2 > 0 \\ \frac{\pi i}{2} H_0^{(1)}(M_{B_s} \sqrt{|F_{(2)}^2|} b_1), & \text{for } F_{(2)}^2 < 0 \end{array} \right), \quad (\text{A4})$$

$$h_f^2(x_1, x_2, x_3, b_1, b_2) = \left\{ \theta(b_1 - b_2) K_0(i\sqrt{x_2 x_3} b_1 M_{B_s}) I_0(i\sqrt{x_2 x_3} b_2 M_{B_s}) \right. \\ \left. + (b_1 \leftrightarrow b_2) \right\} \cdot \left(\begin{array}{ll} K_0(M_{B_s} F_{(3)} b_1), & \text{for } F_{(3)}^2 > 0 \\ \frac{\pi i}{2} H_0^{(1)}(M_{B_s} \sqrt{|F_{(3)}^2|} b_1), & \text{for } F_{(3)}^2 < 0 \end{array} \right), \quad (\text{A5})$$

where J_0 is the Bessel function, K_0 and I_0 are modified Bessel functions $K_0(-ix) = -(\pi/2)Y_0(x) + i(\pi/2)J_0(x)$; $H_0^{(1)}(z)$ is the Hankel function, $H_0^{(1)}(z) = J_0(z) + iY_0(z)$, and $F_{(j)}$'s are defined by

$$F_{(1)}^2 = (x_1 - x_2)x_3, \quad (\text{A6})$$

$$F_{(2)}^2 = (x_1 - x_2)x_3, \quad (\text{A7})$$

$$F_{(3)}^2 = x_1 + x_2 + x_3 - x_1 x_3 - x_2 x_3. \quad (\text{A8})$$

The threshold resummation form factor $S_t(x_i)$ is adopted from Ref.[17]

$$S_t(x) = \frac{2^{1+2c}\Gamma(3/2+c)}{\sqrt{\pi}\Gamma(1+c)}[x(1-x)]^c, \quad (\text{A9})$$

where the parameter $c = 0.3$. This function is normalized to unity. More information about the threshold resummation can be found in reference [13, 15].

The Sudakov factors used in the text are defined as

$$S_{ab}(t) = s\left(x_1 m_{B_s}/\sqrt{2}, b_1\right) + s\left(x_3 m_{B_s}/\sqrt{2}, b_3\right) + s\left((1-x_3)m_{B_s}/\sqrt{2}, b_3\right) \\ - \frac{1}{\beta_1} \left[\ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_3\Lambda)} \right], \quad (\text{A10})$$

$$S_{cd}(t) = s\left(x_1 m_{B_s}/\sqrt{2}, b_1\right) + s\left(x_2 m_{B_s}/\sqrt{2}, b_2\right) + s\left((1-x_2)m_{B_s}/\sqrt{2}, b_2\right) \\ + s\left(x_3 m_{B_s}/\sqrt{2}, b_1\right) + s\left((1-x_3)m_{B_s}/\sqrt{2}, b_1\right) \\ - \frac{1}{\beta_1} \left[2 \ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_2\Lambda)} \right], \quad (\text{A11})$$

$$S_{ef}(t) = s\left(x_1 m_{B_s}/\sqrt{2}, b_1\right) + s\left(x_2 m_{B_s}/\sqrt{2}, b_2\right) + s\left((1-x_2)m_{B_s}/\sqrt{2}, b_2\right) \\ + s\left(x_3 m_{B_s}/\sqrt{2}, b_2\right) + s\left((1-x_3)m_{B_s}/\sqrt{2}, b_2\right) \\ - \frac{1}{\beta_1} \left[\ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + 2 \ln \frac{\ln(t/\Lambda)}{-\ln(b_2\Lambda)} \right], \quad (\text{A12})$$

$$S_{gh}(t) = s\left(x_2 m_{B_s}/\sqrt{2}, b_2\right) + s\left(x_3 m_{B_s}/\sqrt{2}, b_3\right) + s\left((1-x_2)m_{B_s}/\sqrt{2}, b_2\right) \\ + s\left((1-x_3)m_{B_s}/\sqrt{2}, b_3\right) - \frac{1}{\beta_1} \left[\ln \frac{\ln(t/\Lambda)}{-\ln(b_3\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_2\Lambda)} \right], \quad (\text{A13})$$

where the function S_{B_s} , S_{ρ^0} , $S_{\eta^{(\prime)}}$ used in the amplitudes are defined as:

$$S_{B_s}(t) = s(x_1 P_1^+, b_1) + 2 \int_{1/b_1}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma(\alpha_s(\bar{\mu})), \quad (\text{A14})$$

$$S_{\rho^0}(t) = s(x_2 P_2^+, b_2) + s((1-x_2) P_2^+, b_2) + 2 \int_{1/b_2}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma(\alpha_s(\bar{\mu})), \quad (\text{A15})$$

$$S_{\eta^{(\prime)}}(t) = s(x_3 P_3^-, b_3) + s((1-x_3) P_3^-, b_3) + 2 \int_{1/b_3}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma(\alpha_s(\bar{\mu})). \quad (\text{A16})$$

where the so called Sudakov factor $s(Q, b)$ resulting from the resummation of double logarithms is given as:

$$s(Q, b) = \int_{1/b}^Q \frac{d\mu}{\mu} \left[\ln \left(\frac{Q}{\mu} \right) A(\alpha_s(\bar{\mu})) + B(\alpha_s(\bar{\mu})) \right] \quad (\text{A17})$$

with

$$A = C_F \frac{\alpha_s}{\pi} + \left[\frac{67}{9} - \frac{\pi^2}{3} - \frac{10}{27} n_f + \frac{2}{3} \beta_0 \ln \left(\frac{e^{\gamma_E}}{2} \right) \right] \left(\frac{\alpha_s}{\pi} \right)^2, \quad (\text{A18})$$

$$B = \frac{2}{3} \frac{\alpha_s}{\pi} \ln \left(\frac{e^{2\gamma_E-1}}{2} \right). \quad (\text{A19})$$

Here $\gamma_E = 0.57722 \dots$ is the Euler constant, n_f is the active quark flavor number. For the detailed derivation of the Sudakov factors, see Ref.[34]. The hard scale t_i 's in the above equations are chosen as:

$$t_e^1 = \max(\sqrt{x_3} m_{B_s}, 1/b_1, 1/b_3), \quad (\text{A20})$$

$$t_e^2 = \max(\sqrt{x_1} m_{B_s}, 1/b_1, 1/b_3), \quad (\text{A21})$$

$$t_e^3 = \max(\sqrt{x_3} m_{B_s}, 1/b_2, 1/b_3), \quad (\text{A22})$$

$$t_e^4 = \max(\sqrt{x_2} m_{B_s}, 1/b_2, 1/b_3), \quad (\text{A23})$$

$$t_f = \max(\sqrt{x_1 x_3} m_{B_s}, \sqrt{x_2 x_3} m_{B_s}, 1/b_1, 1/b_2), \quad (\text{A24})$$

$$t_f^1 = \max(\sqrt{x_2 x_3} m_{B_s}, 1/b_1, 1/b_2), \quad (\text{A25})$$

$$t_f^2 = \max(\sqrt{x_1 + x_2 + x_3 - x_1 x_3 - x_2 x_3} m_{B_s}, \sqrt{x_2 x_3} m_{B_s}, 1/b_1, 1/b_2). \quad (\text{A26})$$

They are given as the maximum energy scale appearing in each diagram to kill the large logarithmic radiative corrections.

-
- [1] P. Ball et al., hep-ph/0003238; S. Stone, talk given at the Aspen Winter Conference, Aspen, Colorado, USA, Feb.12-18, 2006; R. Fleischer, Phys. Rep. **370**, 537 (2002); T. Hurth, Rev. Mod. Phys. **75**, 1159(2003).
 - [2] A. Ali, G. Kramer and C.D. Lü, Phys. Rev. D **58**, 094009 (1998); *ibid.* **59**, 014005 (1999).
 - [3] Y.-H. Chen, H.-Y. Cheng, B. Tseng, and K.-C. Yang, Phys. Rev. D **60**, 094014 (1999); H.-Y. Cheng and K.-C. Yang, Phys. Rev. D **62**, 054029 (2000).
 - [4] M. Beneke, G. Buchalla, M. Neubert, and C.T. Sachrajda, Phys. Rev. Lett. **83**, 1914 (1999); Nucl. Phys. B **591**, 313 (2000);
 - [5] Y.H. Chen, H.Y. Cheng, B. Tseng, Phys. Rev. D **59**, 074003 (1999).
 - [6] B. Tseng, Phys. Lett. B **446**, 125 (1999).
 - [7] J.F. Sun, G.H. Zhu, and D.S. Du, Phys. Rev. D **68**, 054003 (2003).
 - [8] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003).
 - [9] D. Zhang, Z.J. Xiao, and C.S. Li, Phys. Rev. D **64**, 014014 (2001).
 - [10] J.D. Bjorken, Nucl. Phys. B (Proc. Suppl.) **11**, 325 (1989).
 - [11] G.P. Lepage and S.J. Brodsky, Phys. Rev. D **22**, 2157 (1980); J. Potts and G. Sterman, Nucl. Phys. B **225**, 62 (1989).
 - [12] H.N. Li and B. Tseng, Phys. Rev. D **57**, 443 (1998).
 - [13] H.N. Li and H.-L. Yu, Phys. Rev. D **53**, 2480(1996); Y.-Y. Keum, H.N. Li, A.I. Sanda, Phys. Rev. D **63**, 054008(2001).
 - [14] C.D. Lu, K. Ukai, and M.Z. Yang, Phys. Rev. D **63**, 074009(2001).
 - [15] H.N. Li, Phys. Rev. D **66**, 094010(2002); H.N. Li and K. Ukai, Phys. Lett. B **555**, 197(2003).
 - [16] G. Buchalla, A.J. Buras, M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
 - [17] T. Kurimoto, H.N. Li, and A.I. Sanda, Phys. Rev. D **65**, 014007 (2002); C.D. Lu and M.Z. Yang, Eur. Phys. J. C **28**, 515 (2003).
 - [18] E. Kou and A.I. Sanda, Phys. Lett. B **525**, 240 (2002).
 - [19] Particle Data Group, W.-M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
 - [20] T. Feldmann, P. Kroll and B. Stech, Phys. Rev. D **58**, 114006 (1998); Phys. Lett. B **449**, 339 (1999); T. Feldmann, Int. J. Mod. Phys. A **15**, 159 (2000).
 - [21] R. Escribano and J.M. Frere, JHEP 0506 (2005) 029.
 - [22] Y.Y. Charng, T. Kurimoto, and H.N. Li, Phys. Rev. D **74**, 074024 (2006).
 - [23] X. Liu, H.S. Wang, Z.J. Xiao, L.B. Guo, and C.D. Lu, Phys. Rev. D **73**, 074002(2006).
 - [24] H.S. Wang, X. Liu, Z.J. Xiao, L.B. Guo, and C.D. Lu, Nucl. Phys. B **738**, 243(2006).
 - [25] Z.J. Xiao, X.F. Chen and D.Q. Guo, hep-ph/0608222, Eur. Phys. J. C **49** (2007) in press.
 - [26] Z.J. Xiao, X. Liu, and H.S. Wang, hep-ph/0606177.
 - [27] C.D. Lü, K. Ukai, and M.Z. Yang, Phys. Rev. D **63**, 074009 (2001).
 - [28] CDF Collaboration, D. Acosta et al., Phys. Rev. Lett. **95**, 031801(2005).
 - [29] P. Ball, V.M. Braun, Y. Koike, and K. Tanaka, Nucl. Phys. B **529**, 323 (1998).
 - [30] P. Ball, J. High Energy Phys. 9809, 005 (1998); *ibid* , 9901, 010 (1999).
 - [31] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014015 (2005).
 - [32] E. Kou, Phys. Rev. D **63**,054027(2001).
 - [33] Z.J. Xiao, D.Q. Guo and X.F. Chen, hep-ph/0607219, Phys. Rev. D **75** (2007) in press.
 - [34] H.N. Li and H.L. Yu, Phys. Rev. Lett. **74**, 4388 (1995); Phys. Lett. B **353**, 301 (1995).